



Coffin Siris Syndrome

First description and alternative names

The Coffin Siris syndrome was first described by Coffin and Siris in 1970. The researchers described three affected girls with intellectual disability, postnatal growth retardation, lax joints, brachydactyly of the fifth digits with absent nails, dislocation of radial heads, and absence of the fifth terminal phalanges (Coffin & Siris 1970). Alternative names include “Dwarfism- Onychodysplasia”, “Short Stature-Onychodysplasia”, “Fifth Digit syndrome”, and “Mental Retardation and Hypoplastic 5th Fingernails”.

Genetics and molecular biology

Coffin-Siris syndrome is now regarded as one of the BAF-1 disorders (Mannino et al. 2018). It is now regarded as of equal sex distribution in 60 molecularly confirmed cases (Santen et al. 2014). An autosomal dominant inheritance pattern with complete penetrance is suggested (Schrier-Vergano et al. 2018).

Studies have examined the candidate region for Coffin Siris Syndrome. An infant with findings consistent with Coffin Siris syndrome was reported to have an apparently balanced translocation with breakpoints at 1q21.3 and 7q34 (Mcperson et al. 1997). Other research suggested a candidate region for Coffin Siris at 7q32->34 (McGhee et al. 2000). A partial trisomy 9p gene change has also resulted in a Coffin Siris phenotype (Temtamy et al. 2007). Recent advances in molecular genetics such as whole-exome sequencing has seen the identification of SMARCE1 and another seven genes (SMARCB1, SMARCA4, SMARCA2, ARID1A, ARID1B, SOX11 and PHF6) as being implicated in the syndrome (Schrier-Vergano et al. 2018).

Incidence/prevalence

Approximately 200 cases of Coffin Siris syndrome have been reported as of 2018 (Mannino et al. 2018).

Physical features and natural history

Classic clinical criteria for the diagnosis of Coffin Siris syndrome include developmental delay, hirsutism, coarse facies, feeding problems, and hypoplasia or aplasia of the distal fifth digits and nails. Mannino et al. (2018) stressed the importance of molecular testing to confirm the diagnosis, giving an example of a patient with genetically confirmed CSS who had normal 5th digit fingers and toes bilaterally. Additional characteristics of Coffin Siris syndrome include feeding or sucking difficulties, wide mouth, thick lips, hypotonia, flat nasal bridge with broad nose, sparse scalp hair, thick and laterally placed eyebrows, microcephaly, abnormal ear placement, abnormal or delayed dentition, high arched palate, delayed bone age, coxa valga, frequent otitis media, and respiratory infections (Fleck et al. 2001). Head circumference-for-age percentile is generally reported to be less than the 3% to 10%. There are several reports of individuals with Dandy-Walker malformations or Dandy-Walker variants. Seizures are infrequently reported.

Behavioral and psychiatric characteristics

A few individuals have been described as aggressive or self-injurious while some have been characterized as having happy personalities.

Neuropsychological characteristics

The level of developmental delay varies from mild to moderate; the syndrome was initially characterized with having significant developmental delays (Brautbar et al. 2008). Expressive speech is significantly delayed while receptive speech may be slightly less impacted. Motor skills are somewhat less delayed. Today, individuals are generally reported to attend Special Education classes with an IEP (individualized education plan).

Available guidelines for behavioral assessment/treatment/management

Frequent infections have been reported for individuals diagnosed with Coffin Siris syndrome. Respiratory infections may be related to hypotonia, poor sucking, and aspiration besides increased susceptibility. Consultation with feeding clinics, G tube placement, fundoplication, thickened infant feedings, and careful positioning post-feeding may be helpful. Early cardiac evaluation is suggested. Indications for surgical or medical treatment of congenital heart disease are managed the same as the general population. CNS imaging and GI evaluation may be considered. Renal ultrasound evaluation is suggested to investigate for infrequently reported renal abnormalities. Hernias and tear duct occlusion are treated as medically indicated. Myringotomy and adenoidectomy when indicated may decrease recurrent otitis. Regular audiological screening for hearing loss is recommended because of frequent otitis media. Ophthalmologic evaluations are suggested because of reported vision problems. Pediatric dental evaluation is appropriate; primary teeth are potentially retained long term.

Useful Websites

NIH, Office of Rare Diseases Research: rarediseases.info.nih.gov/

References

- 1. Brautbar A., Ragsdale J. & Shinawi M. (2008)** Is this Coffin-Siris Syndrome or the BOD Syndrome? *Am J Med Genet* 149, 559-62.
- 2. Coffin G.S. & Siris E. (1970)** Mental Retardation With Absent Fifth Fingernail and Terminal Phalanx. *Amer J Dis Child* 119, 433-9.
- 3. Fleck B.J., Pandya A., Vanner L., Kerkering K. & Bodurtha J. (2001)** Coffin Siris Syndrome: Review and Presentation of New Cases From a Questionnaire Study. *Am J Med Genet* 99, 1-7.
- 4. Mannino E., Miyawaki H., Santen G., Schrier-Vergano S. (2018).** First data from a parent-reported registry of 81 individuals with Coffin–Siris syndrome: Natural history and management recommendations. *Am J Med Genet A.* 176(11): 2250-2258.
- 5. McGhee E.M., Klump C.J., Bitts S.M., Cotter P.D. & Lammer E.J. (2000)** *Am J Med Genet* 93, 241- 3.
- 6. McPherson E.W., Laneri G., Clemens M.M., Kochmar S.J. & Surti U. (1997)** Apparently balanced t(1;7)(q21.3;q34) in an Infant With Coffin-Siris syndrome. *Am J Med Genet* 71, 430-3.
- 7. Santen G., Clayton-Smith J., ARD1B-CSS Consortium. (2014).** The ARID1B Phenotype: What we have learned so far. *Am J Med Genet C Semin Med Genet.* 166C: 276–89.

8. **Schrier-Vergano S, Santen G, Wieczorek D, Wollnik B, Matsumoto N, Deardorff M. (2018).** Coffin-Siris Syndrome. In: Adam MP, Ardinger HH, Pagon RA, et al., editors. GeneReviews® [Internet]. Seattle (WA): University of Washington, Seattle; 1993-2019.
9. **Temtamy S.A., Kamel A.K., Ismail S., Helmy N.A., Aglan M.S., El Gammal M., El Ruby M. & Mohamed A.M. (2007)** Phenotypic and cytogenetic spectrum of 9p trisomy. *Genet Couns* 18(1), 29- 48.
10. **Tsurusaki Y; Okamoto N; Ohashi H; Mizuno S; Matsumoto N; Makita Y; Fukuda M; Isidor B; Perrier J; Aggarwal S; Dalal AB; Al-Kindy A; Liebelt J; Mowat D; Nakashima M; Saito H; Miyake N; Matsumoto N.** Coffin-Siris syndrome is a SWI/SNF complex disorder. *Clinical Genetics*. 85(6):548-54, 2014 Jun.

Judith Hiemenga, Srinivasan Sathyanarayanan & Joann Bodurtha, 2010

Revised Navin Dadlani & Stewart Einfeld, June 2019

Copyright © 2019

The SSBP hopes that readers will find the syndrome information sheets useful. They represent the views of the authors who kindly prepared them, and not necessarily those of the SSBP.